

Economic Burden of Macrophage Activation Syndrome (MAS) in Patients With Still’s Disease (Systemic Juvenile Idiopathic Arthritis [sJIA] and Adult-onset Still’s Disease [AOSD]): Analysis of a US National Administrative Claims Database

Michael Marrone, PhD, MPH; [Christopher McPherson](#), PharmD; Abiola Oladapo, PhD
Sobi, Inc., Waltham, MA, USA

CONCLUSIONS

- **Patients with macrophage activation syndrome (MAS) associated with Still’s disease incurred significantly higher healthcare resource utilization (HCRU) and related costs compared with patients with Still’s disease without MAS**
- **Novel MAS-targeted treatments may alleviate the overall disease and economic burden on patients and the healthcare system**

BACKGROUND

- MAS is a rare, potentially-fatal, systemic hyperinflammatory syndrome that occurs as a complication of rheumatologic disease, particularly Still’s disease (systemic juvenile idiopathic arthritis [sJIA] and adult-onset Still’s disease [AOSD])^{1,2}
- Untreated or inadequately treated MAS may rapidly progress to multiorgan failure and death³
- There is only 1 recently approved therapy for MAS in Still’s disease, emapalumab; however; the economic burden (HCRU and costs) of MAS in patients with Still’s disease in the United States (US) is unknown⁴

OBJECTIVE

To assess HCRU and costs associated with MAS in patients with Still’s disease using a US national administrative claims database

METHODS

- In this retrospective analysis, the IQVIA PharMetrics Plus administrative claims database from 2019 to 2024 was used to identify:
 - Patients with Still’s disease (International Classification of Diseases [ICD]-10: M082), including AOSD (ICD-10: M061) and sJIA (ICD-10: M08) with ≥6 months of continuous enrollment prior to Still’s disease diagnosis (index date) and ≥12 months of continuous enrollment post-index without a MAS diagnosis (ICD-10:D76.1)
 - Patients with Still’s disease with ≥6 months of continuous enrollment prior to earliest MAS diagnosis (index date) observed in the study period and ≥12 months of continuous enrollment post-MAS diagnosis
- A propensity score (PS) was used to match patients (1:1) with Still’s disease with and without MAS. Baseline variables used for PS matching were age at Still’s disease diagnosis, sex, race/ethnicity, underlying Still’s disease diagnosis (AOSD or sJIA), and Elixhauser Comorbidity Index (ECI)
- Odds ratios (ORs) were evaluated for binary variables (≥1 visit) using logistic regression
- Generalized linear models were used to estimate the impact of MAS on HCRU (Poisson distribution) and costs (gamma distribution)
- Mean differences in HCRU and costs between the 2 groups were estimated using unadjusted models and models that adjusted for age at Still’s disease diagnosis, underlying Still’s disease diagnosis, sex, and ECI during the baseline period

RESULTS

- After PS matching, 45 patients with Still’s disease and MAS and 45 patients with Still’s disease only were included in the analysis
- In both cohorts, the mean age at diagnosis of Still’s disease was 26.8 years, more than one-half of the patients were female, and more (>60%) patients had AOSD than sJIA (**Table 1**)
- MAS was diagnosed at a mean (standard deviation [SD]) of 4.6 (10.1) years after Still’s disease diagnosis
- The mean (SD) ECI was higher in patients with Still’s disease and MAS than in patients with Still’s disease without MAS (**Table 1**)
- In both cohorts, >40% of the patients had 1 to 2 comorbidities. The most common comorbidities were rheumatoid arthritis and cardiac arrythmia (**Table 1**)

Table 1. Patient demographics and clinical characteristics at Still’s disease diagnosis

	Still’s disease with MAS (n = 45)	Still’s disease without MAS (n = 45)
Age at diagnosis of Still’s disease, mean (SD), years	26.8 (16.0)	26.8 (15.1)
Sex, n (%)		
Female	27 (60.0)	25 (55.6)
Still’s disease diagnosis		
Adult-onset Still’s disease, n (%)	30 (66.7)	31 (68.9)
Systemic juvenile idiopathic arthritis, n (%)	15 (33.3)	14 (31.1)
Elixhauser Comorbidity Index, mean (SD)	2.1 (2.8)	1.6 (1.5)
Elixhauser comorbidities, n (%)		
Rheumatoid arthritis	19 (42.2)	22 (48.9)
Fluid and electrolyte disorder	12 (26.7)	3 (6.7)
Cardiac arrhythmia	10 (22.2)	6 (13.3)
Depression	9 (7.3)	17 (13.7)
Anemia deficiency	3 (6.7)	7 (15.6)
Type of insurance, n (%)		
Commercial insurance	16 (35.6)	21 (46.7)
Self-insured	28 (62.2)	22 (48.9)


MAS, macrophage activation syndrome; SD, standard deviation.

- Most patients in both cohorts were self-insured or had commercial insurance and were covered by a Preferred Provider Organization

All-cause HCRU

- In the 12-month post-index period, patients with Still’s disease and MAS had significantly more HCRU, including inpatient (IP) admissions (OR: 64.1; P<0.001) and/or emergency room (ER) visits (OR: 2.99; P=0.04) than patients without MAS (**Figure 1**)


Figure 1. All-cause HCRU (proportion of patients) during the post-index period



CI, confidence interval; HCRU, healthcare resource utilization; MAS, macrophage activation syndrome; OR, odds ratio.

- Patients with Still’s disease and MAS had significantly more frequent IP admissions (P<0.001), ER visits (P<0.001), outpatient (OP) visits (P<0.001), and longer IP stays (P<0.001) than patients with Still’s disease without MAS (**Figure 2**)
- These results remained robust and did not change when adjusted for age at Still’s disease diagnosis, underlying Still’s disease diagnosis (AOSD, or sJIA), sex, and ECI

Figure 2. All-cause HCRU during the post-index period




*Limited to patients with ≥1 inpatient admission. P-values <0.001 for comparisons across all types of visits. CI, confidence interval; HCRU, healthcare resource utilization; MAS, macrophage activation syndrome.

All-cause costs and rheumatologic-disease associated costs

- Patients with Still’s disease and MAS incurred significantly higher total all-cause (\$318,828 vs \$19,233; P=0.008) and rheumatologic-related (\$163,099 vs \$1930; P=0.02) healthcare costs than patients without MAS (**Figure 3**)
- These results remained robust and did not change when adjusted for age at Still’s disease diagnosis, underlying Still’s disease diagnosis (AOSD, or sJIA), sex, and ECI

Figure 3. All-cause healthcare costs* (US dollars) in the post-index period



*Adjusted to 2024 US dollars using the medical component of the Consumer Price Index.

†Inpatient costs limited to patients with ≥1 inpatient admission. MAS, macrophage activation syndrome.

All-cause HCRU and costs 6 months pre- and post-MAS diagnosis


- Patients with Still’s disease had significantly higher HCRU (**Table 2**) and all-cause healthcare costs after progressing to MAS than when they had Still’s disease only (\$71,919 vs \$5957; **Figure 4**)

Table 2. All-cause HCRU 6 months pre- and post-MAS diagnosis

	Pre-MAS (n = 45)	Post-MAS (n = 45)	P-value
Inpatient admissions, n			
Median (Q1, Q3)	0 (0, 1)	1 (1, 2)	<0.001
Inpatient length of stay,* (days)			
Median (Q1, Q3)	7 (5.5, 11)	9 (5.5, 12)	0.5
Emergency room visits, n			
Median (Q1, Q3)	0 (0, 1)	0 (0, 1)	1.0
Outpatient visits, n			
Median (Q1, Q3)	4 (1, 8)	11 (4, 26)	<0.001
Laboratory visits, n			
Median (Q1, Q3)	0 (0, 2)	1 (0, 2)	0.007
Pharmacy claims, n			
Median (Q1, Q3)	0 (0, 6)	5 (0, 19)	<0.001

*Limited to patients with ≥1 inpatient admission. HCRU, healthcare resource utilization; MAS, macrophage activation syndrome; Q1, first quartile; Q3, third quartile.

Figure 4. All-cause healthcare costs* (US dollars) 6 months pre- and post-MAS diagnosis



*Adjusted to 2024 US dollars using the medical component of the Consumer Price Index.

†Inpatient costs limited to patients with ≥1 inpatient admission. MAS, macrophage activation syndrome.

LIMITATIONS

- As with other retrospective studies, there is a risk of missing or incomplete information
- PS matching is limited to the baseline variables for which data are available

REFERENCES

1. Grom AA, Horne A, De Benedetti F. *Nat Rev Rheumatol*. 2016;12:259–268. 2. Hentzer JJ. *N Engl J Med*. 2025;392:584–598. 3. De Benedetti F, Grom AA, Brogan PA, et al. *Ann Rheum Dis*. 2023;82:857–865. 4. Gamrani*** (emapalumab-kzq) injection, prescribing information. Updated 6/2025. Accessed August 18, 2025.

CONFLICTS OF INTEREST

Dr Marrone, Dr McPherson, and Dr Oladapo are employees of Sobi, Inc.

ACKNOWLEDGMENTS

This study was funded by Swedish Orphan Biovitrum (Sobi Inc.). Editorial and medical writing support was provided by Aparna Nori, PhD, CMPP, of rareLife solutions, Westport, CT, USA, and funded by Sobi, Inc. This poster was created by the authors in accordance with Good Publication Practice (GPP) 2022 guidelines (<https://www.impp.org/gpp-2022>). Sobi reviewed and provided feedback on the poster. The authors had full editorial control of the poster and provided their final approval of all content.